

10/566,413

=> d his

(FILE 'HOME' ENTERED AT 11:00:51 ON 01 OCT 2008)

FILE 'REGISTRY' ENTERED AT 11:01:14 ON 01 OCT 2008
L1 STRUCTURE UPLOADED

FILE 'CASREACT' ENTERED AT 11:01:56 ON 01 OCT 2008
L2 0 S L1
L3 2 S L1 SSS FUL

=> d l3 ibib all

L3 ANSWER 1 OF 2 CASREACT COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 144:292785 CASREACT

TITLE: Process for preparation of 11-[4-[2-(2-hydroxyethoxy)ethyl]-1-piperazinyl]dibenzo[b,f][1,4]thiazepine (Quetiapine) from 2-amino-2'-carboxydiphenyl sulfide and 1-hydroxyethoxyethylpiperazine.

INVENTOR(S): Pathak, Shailendra; Sharma, Jitendra; Kaushik, Geetesh; Thaper, Rajesh Kumar; Dubey, Sushil Kumar

PATENT ASSIGNEE(S): Jubilant Organosys Limited, India

SOURCE: PCT Int. Appl., 23 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2006027789	A1	20060316	WO 2004-IN281	20040908
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
RW:	AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			

IN 2006DN04348	A	20070713	IN 2006-DN4348	20060727
----------------	---	----------	----------------	----------

PRIORITY APPLN. INFO.:	WO 2004-IN281	20040908
------------------------	---------------	----------

AN 144:292785 CASREACT

TI Process for preparation of 11-[4-[2-(2-hydroxyethoxy)ethyl]-1-piperazinyl]dibenzo[b,f][1,4]thiazepine (Quetiapine) from 2-amino-2'-carboxydiphenyl sulfide and 1-hydroxyethoxyethylpiperazine.

IN Pathak, Shailendra; Sharma, Jitendra; Kaushik, Geetesh; Thaper, Rajesh Kumar; Dubey, Sushil Kumar

PA Jubilant Organosys Limited, India

SO PCT Int. Appl., 23 pp.

CODEN: PIXXD2

DT Patent

LA English

IC ICM C07D281-02

CC 28-22 (Heterocyclic Compounds (More Than One Hetero Atom))

FAN.CNT 1

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2006027789	A1	20060316	WO 2004-IN281	20040908
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
RW:	AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI,			

CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS,
 MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD,
 RU, TJ, TM

IN 2006DN04348 A 20070713 IN 2006-DN4348 20060727

PRAI WO 2004-IN281 20040908

AB A process for preparation of Quetiapine comprises reaction of
 2-amino-2'-carboxydiphenyl sulfide with a phosphorus halide or oxyhalide
 to give an iminohalide which is treated with 1-
 hydroxyethoxyethylpiperazine.

ST Quetiapine prepn; dibenzothiazepinylpiperazinylethoxyethanol prepn;
 aminocarboxydiphenyl sulfide hydroxyethoxyethylpiperazine reaction

IT Bases, reactions
 RL: RGT (Reagent); RACT (Reactant or reagent)
 (inorg.; preparation of Quetiapine from aminocarboxydiphenyl sulfide and
 1-hydroxyethoxyethylpiperazine)

IT Bases, reactions
 RL: RGT (Reagent); RACT (Reactant or reagent)
 (organic; preparation of Quetiapine from aminocarboxydiphenyl sulfide and
 1-hydroxyethoxyethylpiperazine)

IT Phase transfer catalysts
 (preparation of Quetiapine from aminocarboxydiphenyl sulfide and
 1-hydroxyethoxyethylpiperazine)

IT Bicarbonates
 Carbonates, reactions
 Hydrides
 Hydroxides (inorganic)
 Metal alkoxides
 RL: RGT (Reagent); RACT (Reactant or reagent)
 (preparation of Quetiapine from aminocarboxydiphenyl sulfide and
 1-hydroxyethoxyethylpiperazine)

IT 5747-48-8P 19806-43-0P 329216-67-3P
 RL: IMF (Industrial manufacture); RCT (Reactant); SPN (Synthetic
 preparation); PREP (Preparation); RACT (Reactant or reagent)
 (preparation of Quetiapine from aminocarboxydiphenyl sulfide and
 1-hydroxyethoxyethylpiperazine)

IT 111974-69-7P, Quetiapine 111974-72-2P, Quetiapine hemifumarate
 RL: IMF (Industrial manufacture); SPN (Synthetic preparation); PREP
 (Preparation)
 (preparation of Quetiapine from aminocarboxydiphenyl sulfide and
 1-hydroxyethoxyethylpiperazine)

IT 67-56-1, Methanol, uses 67-63-0, Isopropanol, uses 67-64-1, Acetone,
 uses 67-68-5, Dimethyl sulfoxide, uses 68-12-2, Dmf, uses 80-73-9
 120-94-5, N-Methylpyrrolidine 127-19-5, Dimethylacetamide 141-78-6,
 Ethyl acetate, uses
 RL: NUU (Other use, unclassified); USES (Uses)
 (preparation of Quetiapine from aminocarboxydiphenyl sulfide and
 1-hydroxyethoxyethylpiperazine)

IT 88-73-3, o-Chloronitrobenzene 103-76-4, 1-(2-Hydroxyethyl)piperazine
 107-21-1, Ethylene glycol, reactions 110-85-0, Piperazine, reactions
 147-93-3, 2-Mercaptobenzoic acid 577-19-5, o-Bromonitrobenzene
 609-73-4, o-Iodonitrobenzene 1493-27-2, o-Fluoronitrobenzene
 13349-82-1 54920-98-8
 RL: RCT (Reactant); RACT (Reactant or reagent)
 (preparation of Quetiapine from aminocarboxydiphenyl sulfide and
 1-hydroxyethoxyethylpiperazine)


RE.CNT 1 THERE ARE 1 CITED REFERENCES AVAILABLE FOR THIS RECORD

RE

10/566,413


(1) Ici Americas Inc; EP 0240228 A1 1987 CAPLUS

RX(1) OF 20 A + B ==> C...



RX(1) RCT A 147-93-3, B 88-73-3
 RGT D 584-08-7 K₂CO₃
 PRO C 19806-43-0
 CAT 311-28-4 Bu₄N.I
 SOL 67-56-1 MeOH
 CON 4 - 6 hours, room temperature -> 70 deg C


RX(2) OF 20 ...C ==> G...




RX(2) RCT C 19806-43-0
 RGT H 1333-74-0 H₂
 PRO G 54920-98-8
 CAT 7440-05-3 Pd
 SOL 67-56-1 MeOH
 CON 10 - 15 hours, 30 - 35 deg C, 100 psi

RX(3) OF 20 ...G + J ==> K

10/566,413




G



J

(3) \longrightarrow



K

RX(3) RCT G 54920-98-8

STAGE(1)

SOL 10025-87-3 POC13

CON 5 - 6 hours, room temperature \rightarrow 110 deg C

STAGE(2)

RCT J 13349-82-1

RGT L 497-19-8 Na₂CO₃



SOL 108-88-3 PhMe, 872-50-4 NMEP

CON SUBSTAGE(2) 6 - 8 hours, reflux

PRO K 111974-69-7

RX(4) OF 20 ...G + P \implies Q...

10/566,413



RX(4) RCT G 54920-98-8

STAGE(1)

SOL 10025-87-3 POC13

CON 5 - 6 hours, room temperature -> 110 deg C

STAGE(2)

RCT P 110-85-0

SOL 108-88-3 PhMe



CON SUBSTAGE(1) 110 - 120 deg C

SUBSTAGE(2) 6 - 8 hours

PRO Q 5747-48-8


RX(5) OF 20 ...Q + R ==> K

10/566,413




RX(5) RCT Q 5747-48-8, R 628-89-7
 RGT L 497-19-8 Na₂CO₃
 PRO K 111974-69-7
 SOL 71-36-3 BuOH
 CON 10 - 12 hours, room temperature -> 120 deg C

RX(6) OF 20 ...G + J ==> T...



10/566,413



T

RX(6) RCT G 54920-98-8

STAGE(1)

RGT M 10025-87-3 POC13

SOL 108-88-3 PhMe

CON 5 - 6 hours, room temperature -> 110 deg C

STAGE(2)


RCT J 13349-82-1

SOL 108-88-3 PhMe, 872-50-4 NMEP

CON SUBSTAGE(2) 6 - 8 hours, 110 - 120 deg C

PRO T 329216-67-3

RX(7) OF 20 ...T + U ==> K




T



U

(7) >

10/566,413



K

RX(7) RCT T 329216-67-3

STAGE(1)

RGT V 121-44-8 Et₃N, W 124-63-0 MeSO₂Cl
SOL 75-09-2 CH₂Cl₂
CON SUBSTAGE(3) 2 hours, room temperature

STAGE(2)

RCT U 107-21-1
RGT X 7646-69-7 NaH
SOL 107-21-1 (CH₂OH)₂, 108-88-3 PhMe
CON SUBSTAGE(1) room temperature
SUBSTAGE(2) 10 - 12 hours, room temperature -> 120 deg C

PRO K 111974-69-7

L3 ANSWER 2 OF 2 CASREACT COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 142:240472 CASREACT
 TITLE: Procedure for preparing a pharmaceutically active compound
 INVENTOR(S): Puig Torres, Salvador; Herbera Espinal, Reyes; Dalmases Barjoan, Pere
 PATENT ASSIGNEE(S): Laboratorios Vita, S. A., Spain
 SOURCE: PCT Int. Appl., 22 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2005014590	A2	20050217	WO 2004-IB2527	20040727
WO 2005014590	A3	20050506		
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW RW: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
ES 2223294	A1	20050216	ES 2003-1922	20030808
ES 2223294	B2	20051001		
EP 1660468	A2	20060531	EP 2004-744176	20040727
EP 1660468	B1	20070718		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, PL, SK, HR				
JP 2007501837	T	20070201	JP 2006-523068	20040727
AT 367383	T	20070815	AT 2004-744176	20040727
ES 2290734	T3	20080216	ES 2004-744176	20040727
US 20060189594	A1	20060824	US 2006-566413	20060130
PRIORITY APPLN. INFO.:				
			ES 2003-1922	20030808
			WO 2004-IB2527	20040727

OTHER SOURCE(S): MARPAT 142:240472

AN 142:240472 CASREACT
 TI Procedure for preparing a pharmaceutically active compound
 IN Puig Torres, Salvador; Herbera Espinal, Reyes; Dalmases Barjoan, Pere
 PA Laboratorios Vita, S. A., Spain
 SO PCT Int. Appl., 22 pp.
 CODEN: PIXXD2
 DT Patent
 LA English
 IC ICM C07D417-00
 CC 28-22 (Heterocyclic Compounds (More Than One Hetero Atom))
 FAN.CNT 1

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI WO 2005014590	A2	20050217	WO 2004-IB2527	20040727
WO 2005014590	A3	20050506		

W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW

RW: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG

ES 2223294 A1 20050216 ES 2003-1922 20030808
 ES 2223294 B2 20051001
 EP 1660468 A2 20060531 EP 2004-744176 20040727
 EP 1660468 B1 20070718

R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, PL, SK, HR

JP 2007501837 T 20070201 JP 2006-523068 20040727
 AT 367383 T 20070815 AT 2004-744176 20040727
 ES 2290734 T3 20080216 ES 2004-744176 20040727
 US 20060189594 A1 20060824 US 2006-566413 20060130

PRAI ES 2003-1922 20030808
 WO 2004-IB2527 20040727
 OS MARPAT 142:240472
 GI

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

AB The invention relates to a procedure for preparing quetiapine (I) by reaction between dibenzothiazepine II and a compound P-OCH₂CH₂X [P = alc. protective group resistant to alkaline conditions; especially ethers, e.g., tetrahydropyranyl, CH₂Ph, trityl; X = leaving group, e.g., halogen, mesylate, triflate, nonaflate, tresylate, tosylate, brosylate, nosylate], in the presence of a base, followed by a step of deprotection of ether III and, optionally, obtaining a pharmaceutically acceptable salt thereof. Said procedure permits the obtaining of quetiapine with a high degree of purity under soft temperature conditions, with short reaction times and avoiding the use of toxic solvents.

ST quetiapine prepn; dibenzothiazepine hydroxyethylpiperazino etherification
 IT Hydrolysis
 (acid, of O-protected quetiapine; procedure for preparing quetiapine from a dibenzothiazepine piperazinoethanol derivative)

IT Bases, reactions
 RL: RGT (Reagent); RACT (Reactant or reagent)
 (alkali and alkaline earth metal derivs.; procedure for preparing quetiapine from a dibenzothiazepine piperazinoethanol derivative)



IT Carbonates, reactions
 RL: RGT (Reagent); RACT (Reactant or reagent)
 (alkali metal and alkaline earth derivs.; procedure for preparing quetiapine from a dibenzothiazepine piperazinoethanol derivative)

IT Heterocyclic compounds
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)

- (dibenzothiazepines; procedure for preparing quetiapine from a dibenzothiazepine piperazinoethanol derivative)
- IT Protective groups
(ethers, alkaline resistant; procedure for preparing quetiapine from a dibenzothiazepine piperazinoethanol derivative)
- IT Phase transfer catalysts
(for etherification of a dibenzothiazepine piperazinoethanol derivative; procedure for preparing quetiapine from a dibenzothiazepine piperazinoethanol derivative)
- IT Etherification
(of a dibenzothiazepine piperazinoethanol derivative; procedure for preparing quetiapine from a dibenzothiazepine piperazinoethanol derivative)
- IT Alkali metal hydroxides
Alkaline earth hydroxides
RL: RGT (Reagent); RACT (Reactant or reagent)
(procedure for preparing quetiapine from a dibenzothiazepine piperazinoethanol derivative)
- IT Quaternary ammonium compounds, uses
RL: CAT (Catalyst use); USES (Uses)
(tri-C8-10-alkylmethyl, chlorides, for etherification of a dibenzothiazepine piperazinoethanol derivative; procedure for preparing quetiapine from a dibenzothiazepine piperazinoethanol derivative)
- IT 1310-58-3, Potassium hydroxide, reactions 1310-73-2, Sodium hydroxide, reactions
RL: RGT (Reagent); RACT (Reactant or reagent)
(etherification agent; procedure for preparing quetiapine from a dibenzothiazepine piperazinoethanol derivative)
- IT 1235-23-0, 2-Chloroethyl trityl ether 5631-96-9, 2-(2-Chloroethoxy)-2H-tetrahydropyran 17229-17-3, Benzyl 2-chloroethyl ether 65338-95-6, 2-[(Tetrahydropyran-2-yl)oxy]ethyl p-toluenesulfonate
RL: RCT (Reactant); RACT (Reactant or reagent)
(etherification by, of dibenzothiazepine piperazinoethanol derivative; procedure for preparing quetiapine from a dibenzothiazepine piperazinoethanol derivative)
- IT 329216-67-3, 2-[4-(Dibenzo[b,f][1,4]thiazepin-11-yl)piperazin-1-yl]ethanol
RL: RCT (Reactant); RACT (Reactant or reagent)
(etherification of, with (chloroethoxy)tetrahydropyran and analogs; procedure for preparing quetiapine from a dibenzothiazepine piperazinoethanol derivative)
- IT 311-28-4, Tetrabutylammonium iodide 17455-13-9, 18-Crown-6 32503-27-8, Tetrabutylammonium bisulfate
RL: CAT (Catalyst use); USES (Uses)
(phase-transfer catalyst; procedure for preparing quetiapine from a dibenzothiazepine piperazinoethanol derivative)
- IT 844639-08-3P, 11-[4-[2-(2-Benzyloxyethoxy)ethyl]piperazin-1-yl]dibenzo[b,f][a,4]thiazepine
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
(preparation and acetolysis of; procedure for preparing quetiapine from a dibenzothiazepine piperazinoethanol derivative)
- IT 844639-06-1P, 11-[4-[2-(2-Trityloxyethoxy)ethyl]piperazin-1-yl]dibenzo[b,f][a,4]thiazepine
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
(preparation and acid hydrolysis of; procedure for preparing quetiapine from a

- dibenzothiazepine piperazinoethanol derivative)
- IT 844639-07-2P, 11-[4-[2-(2-Acetoxyethoxy)ethyl]piperazin-1-yl]dibenzo[b,f][a,4]thiazepine
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
 (preparation and basic hydrolysis of; procedure for preparing quetiapine from a
- dibenzothiazepine piperazinoethanol derivative)
- IT 111974-69-7P, Quetiapine
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
 (preparation and reaction of, with fumaric acid; procedure for preparing quetiapine from a dibenzothiazepine piperazinoethanol derivative)
- IT 111974-72-2P, Quetiapine hemifumarate
 RL: SPN (Synthetic preparation); PREP (Preparation)
 (preparation and reaction of, with fumaric acid; procedure for preparing quetiapine from a dibenzothiazepine piperazinoethanol derivative)
- IT 110-17-8, Fumaric acid, reactions
 RL: RCT (Reactant); RACT (Reactant or reagent)
 (procedure for preparing quetiapine from a dibenzothiazepine piperazinoethanol derivative)

RX(1) OF 13 A + B ==> C



C
 YIELD 85%

10/566,413

RX(1)

STAGE(1)

RGT D 1310-73-2 NaOH
SOL 7732-18-5 Water
CON 25 deg C

STAGE(2)

RCT A 329216-67-3
CON 25 deg C

STAGE(3)

RCT B 5631-96-9
CON 25 deg C

STAGE(4)

CAT 32503-27-8 Bu4N.HSO4
CON SUBSTAGE(1) 25 deg C
SUBSTAGE(2) 25 deg C -> 60 deg C
SUBSTAGE(3) 6 hours, 60 deg C
SUBSTAGE(4) 60 deg C -> 20 deg C

STAGE(5)

SOL 7732-18-5 Water, 108-88-3 PhMe
CON 20 - 25 deg C

STAGE(6)


RGT E 7647-01-0 HCl
SOL 7732-18-5 Water
CON 3 hours, 20 - 25 deg C

STAGE(7)


RGT F 584-08-7 K2CO3
SOL 7732-18-5 Water
CON 20 - 25 deg C, pH 10

PRO C 111974-69-7

RX(2) OF 13 A + J ==> K...




A



J

(2) \longrightarrow

10/566,413



K
YIELD 82%

RX(2) RCT A 329216-67-3, J 1235-23-0

STAGE(1)

CON 100 - 110 deg C

STAGE(2)

RGT L 1310-58-3 KOH

CON 45 - 60 minutes, 100 - 110 deg C

STAGE(3)

CAT 17455-13-9 18-Crown-6

CON 2 hours, 100 - 110 deg C

STAGE(4)

SOL 108-88-3 PhMe

CON 100 - 110 deg C

STAGE(5)

SOL 7732-18-5 Water

CON 100 - 110 deg C -> 20 deg C

STAGE(6)

SOL 67-56-1 MeOH, 108-88-3 PhMe

CON SUBSTAGE(1) 35 - 40 deg C

SUBSTAGE(2) 35 - 40 deg C -> 0 deg C

STAGE(7)

SOL 67-56-1 MeOH, 78-93-3 EtCOMe

CON SUBSTAGE(1) reflux

SUBSTAGE(2) reflux -> 20 deg C

SUBSTAGE(3) 1 hour, 20 - 25 deg C


SUBSTAGE(4) 20 - 25 deg C -> 0 deg C

PRO K 844639-06-1


NTE first stage fusion; fifth stage crystn.; last stage recrystn.

RX(3) OF 13 A + P ==> C

10/566,413




A



P

(3) \longrightarrow



C

YIELD 90%

RX(3)

STAGE(1)

RGT D 1310-73-2 NaOH

SOL 7732-18-5 Water

CON 25 deg C

STAGE(2)

RCT A 329216-67-3

CON 25 deg C

STAGE(3)

10/566,413

RCT P 65338-95-6
CON 25 deg C

STAGE(4)

CAT 32503-27-8 Bu4N.HSO4
CON SUBSTAGE(1) 25 deg C
SUBSTAGE(2) 25 deg C -> 60 deg C
SUBSTAGE(3) 8 hours, 60 - 65 deg C
SUBSTAGE(4) 60 - 65 deg C -> 20 deg C

STAGE(5)

SOL 7732-18-5 Water, 108-88-3 PhMe
CON 20 - 25 deg C

STAGE(6)


RGT E 7647-01-0 HCl
SOL 7732-18-5 Water
CON 3 hours, 20 - 25 deg C

STAGE(7)

RGT F 584-08-7 K2CO3
SOL 7732-18-5 Water
CON 20 - 25 deg C, pH 10

PRO C 111974-69-7


RX(4) OF 13 ...Q ==> C



Q

(4) \longrightarrow

10/566,413



C
YIELD 94%

RX(4) RCT Q 844639-07-2

STAGE(1)

SOL 67-56-1 MeOH
CON 20 - 25 deg C

STAGE(2)

RGT L 1310-58-3 KOH
CON 3 hours, 20 - 25 deg C

STAGE(3)


RGT E 7647-01-0 HCl
SOL 7732-18-5 Water
CON 20 - 25 deg C

STAGE(4)

RGT D 1310-73-2 NaOH
SOL 7732-18-5 Water
CON 20 - 25 deg C, pH 10


PRO C 111974-69-7

RX(5) OF 13 ...R + S ==> Q...



10/566,413

(5) \longrightarrow



Q
YIELD 89%

RX(5) RCT R 64-19-7

STAGE(1)

RGT T 10035-10-6 HBr
SOL 64-19-7 AcOH
CON 20 - 25 deg C

STAGE(2)

RCT S 844639-08-3
CON 1.5 hours, 20 - 25 deg C

STAGE(3)

SOL 7732-18-5 Water, 75-09-2 CH₂Cl₂
CON 20 - 25 deg C

STAGE(4)


RGT U 144-55-8 NaHCO₃
CON 20 - 25 deg C

PRO Q 844639-07-2


NTE last stage neutralization

RX(6) OF 13 ...K + W ==> X

10/566,413




K




W

(6) \longrightarrow



X: CM 1



X: CM 2

RX(6) RCT K 844639-06-1

STAGE(1)

CAT 104-15-4 TsOH
SOL 67-56-1 MeOH, 108-88-3 PhMe
CON 4 hours, reflux

STAGE(2)

RGT E 7647-01-0 HCl
SOL 7732-18-5 Water, 108-88-3 PhMe
CON 20 - 25 deg C

STAGE(3)

RGT D 1310-73-2 NaOH
SOL 7732-18-5 Water, 108-88-3 PhMe
CON 20 - 25 deg C, pH 9.5

STAGE(4)

SOL 67-56-1 MeOH

10/566,413

CON 20 - 25 deg C

STAGE(5)

RCT W 110-17-8

CON SUBSTAGE(1) 35 - 45 minutes, 20 - 25 deg C

SUBSTAGE(2) 20 - 25 deg C -> reflux

SUBSTAGE(3) 5 minutes, reflux


SUBSTAGE(4) reflux -> 10 deg C

SUBSTAGE(5) 1 hour, 10 - 15 deg C


PRO X 111974-72-2

NTE (95%;94%)

RX(7) OF 13 A + B ==> C




A



B

(7) \longrightarrow



C

YIELD 82%

RX(7)

STAGE(1)

RGT L 1310-58-3 KOH

SOL 7732-18-5 Water

CON 25 deg C

STAGE(2)

10/566,413

RCT A 329216-67-3
CON 25 deg C

STAGE(3)

RCT B 5631-96-9
CON 25 deg C

STAGE(4)

CAT 17455-13-9 18-Crown-6
CON SUBSTAGE(1) 25 deg C
SUBSTAGE(2) 25 deg C -> 40 deg C
SUBSTAGE(3) 6 hours, 40 deg C
SUBSTAGE(4) 40 deg C -> 20 deg C

STAGE(5)

SOL 7732-18-5 Water, 108-88-3 PhMe
CON 20 - 25 deg C

STAGE(6)


RGT E 7647-01-0 HCl
SOL 7732-18-5 Water
CON 3 hours, 20 - 25 deg C

STAGE(7)


RGT F 584-08-7 K₂CO₃
SOL 7732-18-5 Water
CON 20 - 25 deg C, pH 10

PRO C 111974-69-7

RX(8) OF 13 A + B ==> C




A



B

(8) \longrightarrow

10/566,413



C
YIELD 75%

RX(8)

STAGE(1)

RGT L 1310-58-3 KOH
SOL 7732-18-5 Water
CON 25 deg C

STAGE(2)

RCT A 329216-67-3
CON 25 deg C

STAGE(3)

RCT B 5631-96-9
CON 25 deg C

STAGE(4)

CON SUBSTAGE(1) 25 deg C
SUBSTAGE(2) 25 deg C -> 40 deg C
SUBSTAGE(3) 20 hours, 40 deg C
SUBSTAGE(4) 40 deg C -> 20 deg C

STAGE(5)

SOL 7732-18-5 Water, 108-88-3 PhMe
CON 20 - 25 deg C

STAGE(6)

RGT E 7647-01-0 HCl
SOL 7732-18-5 Water
CON 3 hours, 20 - 25 deg C



STAGE(7)

RGT F 584-08-7 K₂CO₃
SOL 7732-18-5 Water
CON 20 - 25 deg C, pH 10

PRO C 111974-69-7
NTE stage 4 Aliquat 336 catalyst

RX(9) OF 13 A + Z ==> S...

10/566,413



S
YIELD 93%

RX(9)

STAGE(1)

RGT D 1310-73-2 NaOH
SOL 7732-18-5 Water
CON 20 - 25 deg C

STAGE(2)

RCT A 329216-67-3
CON 20 - 25 deg C

STAGE(3)

RCT Z 17229-17-3
CON 20 - 25 deg C

STAGE(4)

CAT 32503-27-8 Bu₄N.HSO₄
CON SUBSTAGE(1) 20 - 25 deg C
SUBSTAGE(2) 20 - 25 deg C -> 60 deg C
SUBSTAGE(3) 9 hours, 60 deg C
SUBSTAGE(4) 60 deg C -> 20 deg C

STAGE(5)

10/566,413

SOL 7732-18-5 Water, 108-88-3 PhMe
CON 20 - 25 deg C

STAGE(6)

RGT E 7647-01-0 HCl
SOL 7732-18-5 Water
CON 5 minutes, 20 - 25 deg C

PRO S 844639-08-3